


Core

Topic/Skill Knowledge:	Definition/Tips	Example
1. Place Value	The <b>value</b> of where a <b>digit</b> is within a number.	In 726, the value of the 2 is 20, as it is in the 'tens' column.
2. Place Value Columns	The names of the columns that <b>determine the value of each digit</b> .  The 'ones' column is also known as the 'units' column.	 <p>PLACE VALUE CHART</p> <p>Millions Hundred Thousands Ten Thousands Thousands Hundreds Tens Ones Decimal Point Tenths Hundredths Thousandths Ten-Thousandths Hundred-Thousandths Millionths</p>
3. Rounding	To make a number simpler but keep its value close to what it was.  If the <b>digit to the right</b> of the rounding digit is <b>less than 5, round down</b> . If the <b>digit to the right</b> of the rounding digit is <b>5 or more, round up</b> .	74 rounded to the nearest ten is 70, because 74 is closer to 70 than 80.  152,879 rounded to the nearest thousand is 153,000.
4. Decimal Place	The <b>position</b> of a digit to the <b>right of a decimal point</b> .	In the number 0.372, the 7 is in the second decimal place.  0.372 rounded to two decimal places is 0.37, because the 2 tells us to round down.  Careful with money - don't write £27.4, instead write £27.40
5. Significant Figure	The significant figures of a number are the digits which <b>carry meaning</b> (ie. are significant) to the size of the number.  The <b>first significant figure</b> of a number <b>cannot be zero</b> .  In a number with a decimal, trailing zeros are not significant.	In the number 0.00821, the first significant figure is the 8.  In the number 2.740, the 0 is not a significant figure.  0.00821 rounded to 2 significant figures is 0.0082.  19357 rounded to 3 significant figures is 19400. We need to include the two zeros at the end to keep the digits in the same place value columns.
6. Truncation	A method of approximating a decimal number by <b>dropping all decimal places</b> past a certain point <b>without rounding</b> .	3.14159265... can be truncated to 3.1415 (note that if it had been rounded, it would become 3.1416)
7. Error Interval	A <b>range of values</b> that a number could have taken before being rounded or truncated.	0.6 has been rounded to 1 decimal place.  The error interval is:

**Core**

**Knowledge:**

	<p>An error interval is written using inequalities, with a <b>lower bound</b> and an <b>upper bound</b>.</p> <p>Note that the lower bound inequality can be 'equal to', but the upper bound cannot be 'equal to'.</p>	$0.55 \leq x < 0.65$ <p>The lower bound is 0.55 The upper bound is 0.65</p>
8. Estimate	To find something <b>close to the correct answer</b> .	An estimate for the height of a man is 1.8 metres.
9. Approximation	<p>When using approximations to estimate the solution to a calculation, <b>round each number in the calculation to 1 significant figure</b>.</p> <p><math>\approx</math> means 'approximately equal to'</p>	$\frac{348 + 692}{0.526} \approx \frac{300 + 700}{0.5} = 2000$ <p>'Note that dividing by 0.5 is the same as multiplying by 2'</p>
10. Rational Number	<p>A number of the form <math>\frac{p}{q}</math>, where <b>p and q are integers</b> and <b>q <math>\neq</math> 0</b>.</p> <p>A number that cannot be written in this form is called an 'irrational' number</p>	<p><math>\frac{4}{9}</math>, 6, <math>-\frac{1}{3}</math>, <math>\sqrt{25}</math> are examples of rational numbers.</p> <p><math>\pi</math>, <math>\sqrt{2}</math> are examples of an irrational numbers.</p>
11. Surd	<p>The <b>irrational number</b> that is a <b>root of a positive integer</b>, whose value cannot be determined exactly.</p> <p>Surds have <b>infinite non-recurring decimals</b>.</p>	<p><math>\sqrt{2}</math> is a surd because it is a root which cannot be determined exactly.</p> <p><math>\sqrt{2} = 1.41421356 \dots</math> which never repeats.</p>
12. Rules of Surds	$\sqrt{ab} = \sqrt{a} \times \sqrt{b}$ $\sqrt{\frac{a}{b}} = \frac{\sqrt{a}}{\sqrt{b}}$ $a\sqrt{c} \pm b\sqrt{c} = (a \pm b)\sqrt{c}$ $\sqrt{a} \times \sqrt{a} = a$	$\sqrt{48} = \sqrt{16} \times \sqrt{3} = 4\sqrt{3}$ $\sqrt{\frac{25}{36}} = \frac{\sqrt{25}}{\sqrt{36}} = \frac{5}{6}$ $2\sqrt{5} + 7\sqrt{5} = 9\sqrt{5}$ $\sqrt{7} \times \sqrt{7} = 7$
13. Rationalise a Denominator	The process of rewriting a fraction so that the <b>denominator contains only rational numbers</b> .	$\frac{\sqrt{3}}{\sqrt{2}} = \frac{\sqrt{3} \times \sqrt{2}}{\sqrt{2} \times \sqrt{2}} = \frac{\sqrt{6}}{2}$

**Core**

**Knowledge:**

		$\frac{6}{3 + \sqrt{7}} = \frac{6(3 - \sqrt{7})}{(3 + \sqrt{7})(3 - \sqrt{7})}$ $= \frac{18 - 6\sqrt{7}}{9 - 7}$ $= \frac{18 - 6\sqrt{7}}{2} = 9 - 3\sqrt{7}$
--	--	---

Links to Pythagoras, area of 2 D shapes, calculations, mixed numbers, geometric sequences, multiplying out brackets, how to leave your answers at the end of questions, using calculations exactly.